



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

A high-resolution imaging X-ray crystal spectrometer for intense laser plasma interaction experiments

H. Chen, M. Bitter, A. Hazi, K. Hill, S. Kerr, E. Magee, S. Nagel, J. Park, M. Schneider, G. Stone, G. Williams, P. Beiersdorfer

February 27, 2014

20th Topical Conference on High-Temperature Plasma
Diagnostics Conference
Atlanta, GA, United States
June 1, 2014 through June 5, 2014

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

A high-resolution imaging X-ray crystal spectrometer for intense laser plasma interaction experiments

Hui Chen¹, M. Bitter², A. Hazi¹, K. W. Hill², S. Kerr³, E. W. Magee¹, S. R. Nagel¹, J. Park¹, M. B. Schneider¹, G. Stone¹, G. J. Williams¹ and P. Beiersdorfer¹

¹*Lawrence Livermore National Laboratory, 7000 East Avenue, Livermore, California 94550-9234, USA*

²*Princeton Plasma Physics Laboratory, Princeton, New Jersey 08543, USA*

³*University of Alberta, Alberta T6G 2R3, Canada*

ABSTRACT:

Precision measurements of electron and ion temperatures through X-ray line broadening can provide critical information on plasma evolution and collisional equilibration processes. Better understanding of electron-ion temperature equilibration in laser produced high energy density (HED) plasmas, including inertial confinement fusion implosions and short-pulse laser heated plasmas, is vital to improving the simulation and prediction of these plasmas. Adapting a technique established for magnetic confinement fusion experiments [1], an imaging crystal spectrometer has been designed and tested for HED plasmas. The instrument uses a spherically bent quartz 211 crystal with radius of 490.8 mm. The test was performed using the Titan laser at Lawrence Livermore National Laboratory to irradiate titanium slabs with laser intensities of 10^{19} - 10^{20} W/cm². He-like and Li-like Ti lines were recorded, from which the spectrometer performance was evaluated. This spectrometer provides very high spectral resolving power ($E/dE > 7000$) while acquiring a one-dimensional image of the source. The characteristics of the spectrometer and some future applications in HED physics are presented.

[1] M. Bitter, *et al.*, *Rev. Sci. Instrum.* **70**, 292 (1999)

This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.